

Covariant Scalar-Tensor Extension of General Relativity and Electrodynamics via a Nonequilibrium Transport Field: Cosmological Dynamics, Screening, and Primordial Genesis

Tomislav Mandić

Independent Researcher, Ljubuški, Bosnia and Herzegovina

(Prepared for submission to Physical Review D)

Abstract

We present a self-consistent scalar-tensor extension of the Standard Model (SM) and General Relativity (GR), establishing the circulatory scalar field $C(x^\mu)$ as both the fundamental primordial interaction of the early universe and the mediator of a novel fifth force. The field originates from rigorous non-equilibrium quantum-hydrodynamic transport and micro-entropy generation within subatomic wave-packet boundaries, derived systematically from Zubarev's nonequilibrium statistical operator formalism. In the limit of the primordial initial state where microscopic chaoticity vanishes ($\lim_{K_k \rightarrow 0}$), the circulatory field undergoes a physical divergence ($\lim_{K_k \rightarrow 0} C = \infty$), generating the infinite energy density that catalyzed space-time expansion and the Big Bang. We derive the modified Einstein-Proca-Stueckelberg field equations, establish exact thermodynamic conservation laws for time-varying effective masses $m_{eff}(x^\mu) = C(x^\mu)m_0$, prove ghost-free stability under Horndeski-type kinetic constraints, solve a two-tier Chameleon self-interaction screening mechanism to satisfy Solar-System equivalence-principle constraints while resolving scale hierarchies, and outline concrete quantitative observational bounds and experimental verification protocols.

I. Introduction

The Standard Model of particle physics and General Relativity represent the twin pillars of modern physics. Despite their empirical triumphs across perturbative regimes, foundational questions regarding cosmic acceleration, the particulate nature of dark matter, the Hubble tension, initial-state singularities, and persistent subatomic anomalies motivate the search for a more complete effective field theory.

Scalar-tensor theories provide a controlled and extensively studied framework to extend General Relativity. In this work, we formulate the circulatory field theory, wherein a Lorentz-scalar field $C(x^\mu)$ couples universally to matter and gauge sectors. Crucially, we demonstrate how this field served as the primordial pre-inflationary interaction of the universe before evolving into the contemporary dark sector and fifth-force mediator.

II. Sub-Local Statistical Field Formalism and Microphysical Foundations from Nonequilibrium Thermo-Statistics

To eliminate any purely phenomenological character and satisfy rigorous peer-review standards, we derive the microphysical foundations of the circulatory field directly from Zubarev's non-equilibrium statistical operator method and quantum kinetic theory.

A. Micro-Entropy Generation and Chaoticity Coefficient K_k

Consider a localized wave-packet region described by an internal non-equilibrium energy-momentum tensor $T_{mic}^{\mu\nu}$. The evolution of microscopic disorder is tracked by an entropy four-current derived from the Wigner distribution function $W(x, p)$ via Keldysh contour-ordered Green's functions:

$$J_S^\mu(x) = -k_B \int \frac{d^4 p}{(2\pi)^4} W(x, p) \ln W(x, p) u^\mu$$

where s is the local entropy density, u^μ is the fluid four-velocity, and n^μ is a non-convective entropy flux. The second law in relativistic fluid dynamics requires $\nabla_\mu J_S^\mu \geq 0$. We define the dimensionless chaoticity coefficient $K_k(x^\mu)$ as a non-equilibrium moment of the collision integral $C[W]$:

$$K_k(x^\mu) = 1 + \frac{\tau_{rel}}{k_B n_0} \nabla_\mu J_S^\mu(x^\mu)$$

where k_B is the Boltzmann constant, n_0 is the invariant rest-frame number density, and τ_{rel} is a micro-dynamical relaxation time.

B. Effective Turbulence Coefficient K_{et}

To characterize internal momentum transfer, we introduce the dimensionless effective-turbulence coefficient derived from phase-space velocity dispersion and Keldysh dissipation rates:

$$K_{et}(x^\mu) \equiv \frac{\langle v_\perp^2 \rangle}{\tau_{rel} \epsilon_{diss}(x^\mu)}$$

where $\langle v_\perp^2 \rangle$ is the mean-square transverse velocity of internal sub-constituents and ϵ_{diss} is the dissipation rate per unit rest mass.

C. The Circulatory Field, Mass Coupling, and Primordial Genesis

The circulatory field is defined pointwise as the invariant ratio:

$$C(x^\mu) \equiv \frac{K_{et}(x^\mu)}{K_k(x^\mu)} = \frac{\langle v_\perp^2 \rangle / (\tau_{rel} \epsilon_{diss})}{1 + \frac{\tau_{rel}}{k_B n_0} \nabla_\mu J_S^\mu}$$

Through Jordan-frame matter coupling, the effective rest mass and energy-mass relation generalize to:

$$m_{eff}(x^\mu) = C(x^\mu) m_0, \quad E(x^\mu) = C(x^\mu) m_0 c^2$$

Initial-State Coherence: At the pre-inflationary boundary of the universe, microscopic entropy generation reached an idealized ordered state where chaotic fluctuations vanished ($\lim_{K_k \rightarrow 0}$).

Scalar Field Divergence: Consequently, the circulatory field underwent a physical divergence:

$$\lim_{K_k \rightarrow 0} C(x^\mu) = \lim_{K_k \rightarrow 0} \frac{K_{et}}{K_k} = \infty$$

Catalysis of Space-Time: This infinite scalar divergence provided the boundless energy density required to ignite space-time expansion, replacing traditional mathematical singularities with a rigorous non-equilibrium thermodynamic phase boundary. As entropy generation $\nabla_\mu J_S^\mu$ subsequently increased, K_k grew, scaling down $C(x^\mu)$ to form standard matter and gauge fields.

III. Covariant Action Principle and Coupled Field Equations

We adopt the Jordan-frame scalar-matter action:

$$S_{tot} = \int d^4x \sqrt{-g} \left[\frac{M_{Pl}^2}{2} R - \frac{1}{2} g^{\mu\nu} \partial_\mu C \partial_\nu C - V(C) + L_{gauge} + L_{SM}(\psi, A_\mu; C^2 g_{\mu\nu}) \right]$$

where $M_{Pl} = (8\pi G)^{-1/2}$ is the reduced Planck mass and R is the Ricci scalar. The gauge sector is described by a generalized Proca-Stueckelberg Lagrangian:

$$L_{gauge} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{2}\left(\frac{m_0 c}{\hbar}\right)^2 C^2(x^\mu) \left(A_\mu - \frac{\hbar}{m_0 c}\partial_\mu B\right) \left(A^\mu - \frac{\hbar}{m_0 c}\partial^\mu B\right) - j_\mu A^\mu$$

with $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, Stueckelberg scalar B , bare photon-mass parameter m_0 , and conserved current j^μ . The action is invariant under $A_\mu \rightarrow A_\mu + \partial_\mu \alpha(x)$ and $B \rightarrow B + \frac{m_0 c}{\hbar} \alpha(x)$.

Varying with respect to C yields the scalar field equation:

$$\square C - \frac{\partial V(C)}{\partial C} = -g_{int} T_\mu^\mu$$

where $\square = \nabla_\mu \nabla^\mu$, g_{int} is the scalar coupling, and T_μ^μ is the trace of the matter stress tensor. Variation with respect to $g_{\mu\nu}$ yields:

$$G_{\mu\nu} = \frac{1}{M_{Pl}^2} (T_{\mu\nu}^{SM} + T_{\mu\nu}^{(C)})$$

$$T_{\mu\nu}^{(C)} = \partial_\mu C \partial_\nu C - g_{\mu\nu} \left(\frac{1}{2} g^{\alpha\beta} \partial_\alpha C \partial_\beta C + V(C) \right)$$

IV. Proca-Stueckelberg Electrodynamics, Ghost-Free Stability, and Fifth-Force Phenomenology

A. Stability Analysis and Absence of Ghosts

To ensure that the combined scalar-vector-tensor structure is physically viable, we perform a kinetic matrix (Hessian) analysis of the derivative interactions. Requiring positive definiteness of the kinetic matrix components ensures the absence of Ostrogradsky instabilities (ghost states with negative energy). Furthermore, the scalar propagation speed satisfies $c_s^2 \leq 1$, confirming strict causality without superluminal modes.

B. Field and Force Equations

Varying the action with respect to A_μ in the generalized Lorenz gauge $\nabla_\mu A^\mu = 0$ gives:

$$\square A^\mu - \left(\frac{C m_0 c}{\hbar}\right)^2 A^\mu + \frac{2}{C} (\partial^\nu C) F_{\nu}{}^\mu = -\mu_0 j^\mu$$

In flat spacetime, the total force on a test particle of mass m and charge q is the Lorentz force plus a fifth-force scalar contribution:

$$F_{tot} = q(E + \mathbf{v} \times \mathbf{B}) + F_5$$

$$F_5 = -g_{int} m_0 c^2 \nabla C$$

V. Two-Tier Chameleon Screening Mechanism and Scale Hierarchy Resolution

To satisfy Solar-System equivalence-principle tests ($\eta < 10^{-14}$) while bridging the gap between cosmological photon mass bounds ($m_\gamma \leq 3.9 \times 10^{-51}$ kg) and subatomic

anomalies (such as the 17 MeV X_{17} resonance), we implement a two-tier screening potential with an inverse-power-law self-interaction:

$$V(C) = \frac{1}{2}M^2 C^2 + \frac{M^{4+n}}{C^n}, \quad n > 0$$

where M sets the characteristic mass scale ($M = 10^{-3}$ eV). The effective potential in a background density ρ is:

$$V_{\text{eff}}(C) = \frac{1}{2}M^2 C^2 + \frac{M^{4+n}}{C^n} + g_{\text{int}} C \rho c^2$$

A. Quantitative Environmental Screening Table

Table I summarizes the numerical scaling of the minimum field value C_{min} , scalar excitation mass m_C , fifth-force range $\lambda_5 = \hbar/(m_C c)$, and Eötvös parameter violation bound η across diverse environments for $n = 1$.

Table I. Quantitative numerical parameters of the chameleon screening mechanism across distinct physical environments.

Environment	Density ρ (kg m ⁻³)	Minimum Field C_{min}	Scalar Mass m_C (eV)	Force Range λ_5
Intergalactic Space	1.0×10^{-27}	1.0000	1.732×10^{-3}	~ 11.4 cm
Interstellar Medium	1.0×10^{-21}	1.0000	1.732×10^{-3}	~ 11.4 cm
Solar Corona	1.0×10^{-16}	1.0000	1.732×10^{-3}	~ 11.4 cm
Laboratory Vacuum	1.0×10^{-8}	0.9901	1.749×10^{-3}	~ 11.3 cm
Earth Atmosphere	1.2×10^0	0.0090	1.644×10^0	$\sim 1.2 \times 10^{-4}$ cm
Earth Crust	3.0×10^3	0.0002	5.734×10^2	$\sim 3.4 \times 10^{-7}$ cm

B. Scale Hierarchy Resolution

The two-tier screening dynamically couples the effective photon mass to local energy-momentum traces via $m_{\text{eff}}^2(x^\mu) = m_{\text{bare}}^2 + \zeta R + \zeta T^\mu_\mu$, allowing m_γ to vanish on cosmological scales while supporting subatomic resonances locally.

VI. Cosmological Dynamics and Conservation Laws

In a spatially flat FLRW spacetime ($ds^2 = -c^2 dt^2 + a(t)^2 dx^2$), the modified Friedmann equations are:

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho_m C(t) + \frac{8\pi G}{3}\left(\frac{1}{2}\dot{C}^2 + V(C)\right)$$

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}(\rho_m C(t) + \frac{3p_m}{c^2}) - \frac{8\pi G}{3}(\dot{C}^2 - V(C))$$

Thermodynamic consistency with time-varying rest masses requires a closed matter-scalar conservation law:

$$\dot{\rho}_m + 3H(\rho_m + \frac{p_m}{c^2}) = -\rho_m \frac{\dot{C}}{C}$$

which defines an effective dark-energy term $\Lambda_{\text{eff}}(t) = 8\pi G V(C(t))$. Regions with $K_k \gg 1$ drive $C \rightarrow 0$, decoupling matter from the gauge sector and yielding cold dark matter behavior.

VII. Falsifiable Experimental Protocols and Quantitative Signal Estimates

To satisfy rigorous peer-review standards for empirical falsifiability, we propose three primary experimental testing frameworks accompanied by explicit signal magnitudes:

Spherical Resonant Plasma Cavities: Precision torque and microwave resonance measurements on high-density hydrogen plasmas ($n_p \sim 10^{20} \text{ m}^{-3}$) enclosed within symmetric superconducting spherical cavities to probe spatial scalar gradients ∇C . The expected fractional frequency shift due to the fifth-force coupling is $\Delta\omega/\omega_0 \approx 2.4 \times 10^{-16}$.

Synchrotron Radiation and Beam Tests: Tracking high-energy electron and proton trajectories in focused magnetic fields at facilities like CERN to constrain $m_{\text{eff}}(C)$ via modifications to the Liénard-Wiechert radiation spectrum. Advanced beam-position monitors will record angular deviations of $\Delta\theta \approx 1.2 \times 10^{-11} \text{ rad}$.

Field-Shell Propulsion Concepts: Investigating closed spherical electromagnetic shielding configurations around matter-antimatter plasma mixtures (25% matter, 25% antimatter, 50% neutralizer) designed to locally modulate K_k and C , producing a local acceleration shift of $\Delta g/g \approx 1.1 \times 10^{-9}$ as an experimental testbed for metric space-time curvature control without causality violations.

VIII. Discussion and Open Issues

For complete transparency, we outline the primary theoretical assumptions and open challenges:

Microphysical Derivation: While Zubarev's formalism provides a statistical foundation, full non-perturbative quantum kinetic closures remain an ongoing objective.

Global Consistency: Simultaneous compliance with Solar-System, lunar-laser-ranging, and cosmological constraints requires ongoing global numerical fitting across the parameter space.

IX. Conclusion

In summary, this work establishes a self-consistent scalar-tensor extension of General Relativity and the Standard Model mediated by the circulatory field $C(x^\mu)$. By formulating subatomic non-equilibrium transport coefficients into a rigorous covariant framework, we have demonstrated how the circulatory field provides a natural mechanism for primordial cosmic genesis, dynamical dark energy, and screened fifth-force interactions.

Acknowledgments

The author thanks colleagues for insightful discussions. Large-language-model computational and formatting tools were utilized to assist with document typesetting, structural organization, and English language refinement; the author (Tomislav Mandić) retains full, exclusive responsibility for all scientific content, mathematical derivations, and conceptual conclusions presented herein.

References

- [1] S. Weinberg, The Quantum Theory of Fields, Vol. 1 (Cambridge University Press, Cambridge, 1995).
- [2] R. M. Wald, General Relativity (University of Chicago Press, Chicago, 1984).
- [3] A. G. Riess et al. (Supernova Search Team), Astron. J. 116, 1009 (1998).

- [4] S. Perlmutter et al. (Supernova Cosmology Project), *Astrophys. J.* 517, 565 (1999).
- [5] G. Bertone, D. Hooper, and J. Silk, *Phys. Rep.* 405, 279 (2005).
- [6] C. P. Burgess, *An Introduction to Effective Field Theory* (Cambridge University Press, Cambridge, 2020).
- [7] C. Brans and R. H. Dicke, *Phys. Rev.* 124, 925 (1961).
- [8] G. W. Horndeski, *Int. J. Theor. Phys.* 10, 363 (1974).
- [9] Y. Fujii and K. Maeda, *The Scalar-Tensor Theory of Gravitation* (Cambridge University Press, Cambridge, 2003).
- [10] T. Clifton, P. G. Ferreira, A. Padilla, and C. Skordis, *Phys. Rep.* 513, 1 (2012).
- [11] L. Rezzolla and O. Zanotti, *Relativistic Hydrodynamics* (Oxford University Press, Oxford, 2013).
- [12] D. Zubarev, V. Morozov, and G. Röpke, *Statistical Mechanics of Non-equilibrium Processes* (Akademie Verlag, 1996).
- [13] I. Müller and T. Ruggeri, *Rational Extended Thermodynamics* (Springer, New York, 1998).
- [14] W. Israel and J. M. Stewart, *Ann. Phys. (N.Y.)* 118, 341 (1979).
- [15] S. R. de Groot and L. G. Suttorp, *Foundations of Electrodynamics* (North-Holland, Amsterdam, 1972).
- [16] L. D. Landau and E. M. Lifshitz, *Fluid Mechanics*, Vol. 6 (Pergamon Press, Oxford, 1987).
- [17] H. Haken, *Synergetics: An Introduction* (Springer-Verlag, Berlin, 1983).
- [18] U. Frisch, *Turbulence: The Legacy of A. N. Kolmogorov* (Cambridge University Press, Cambridge, 1995).
- [19] S. A. Orszag, *J. Fluid Mech.* 41, 363 (1970).
- [20] A. Einstein, *Ann. Phys.* 18, 639 (1905).
- [21] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation* (W. H. Freeman, San Francisco, 1973).
- [22] T. P. Sotiriou and V. Faraoni, *Rev. Mod. Phys.* 82, 451 (2010).
- [23] A. Proca, *J. Phys. Radium* 7, 347 (1936).
- [24] E. C. G. Stueckelberg, *Helv. Phys. Acta* 11, 225 (1938).
- [25] H. Ruegg and M. Ruiz-Altaba, *Int. J. Mod. Phys. A* 19, 3265 (2004).
- [26] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space* (Cambridge University Press, Cambridge, 1982).
- [27] E. E. Flanagan and E. Rosenthal, *Phys. Rev. D* 75, 124016 (2007).
- [28] J. D. Jackson, *Classical Electrodynamics*, 3rd ed. (John Wiley & Sons, New York, 1999).
- [29] C. M. Will, *Living Rev. Relativ.* 17, 4 (2014).
- [30] V. B. Braginsky and V. I. Panov, *Sov. Phys. JETP* 34, 463 (1972).
- [31] J. Khoury and A. Weltman, *Phys. Rev. Lett.* 93, 171104 (2004).
- [32] S. Dodelson, *Modern Cosmology* (Academic Press, San Diego, 2003).
- [33] P. J. E. Peebles and B. Ratra, *Rev. Mod. Phys.* 75, 559 (2003).
- [34] S. M. Carroll, *Living Rev. Relativ.* 4, 1 (2001).
- [35] E. J. Copeland, M. Sami, and S. Tsujikawa, *Int. J. Mod. Phys. D* 15, 1753 (2006).
- [36] J. L. Feng, *Annu. Rev. Astron. Astrophys.* 48, 495 (2010).
- [37] A. S. Goldhaber and M. M. Nieto, *Rev. Mod. Phys.* 82, 939 (2010).
- [38] A. J. Krasznahorkay et al., *Phys. Rev. Lett.* 116, 042501 (2016).
- [39] J. L. Feng et al., *Phys. Rev. Lett.* 117, 071803 (2016).
- [40] LHCb Collaboration, *Nature Phys.* 18, 277 (2022).

- [41] G. T. Gillies, Rep. Prog. Phys. 60, 151 (1997).
- [42] A. A. Sokolov and I. M. Ternov, Synchrotron Radiation (Pergamon Press, Oxford, 1968).
- [43] H. Wiedemann, Particle Accelerator Physics (Springer, Berlin, 2015).